

Patient Information

YESCARTA™ can cause side effects that can lead to death.

Call or see your oncologist or get emergency help RIGHT AWAY if you have any of these symptoms:

- Fever (100.4°F/38°C or higher)
- Difficulty breathing
- Chills or shaking chills
- Confusion
- Dizziness or lightheadedness
- Severe nausea, vomiting, or diarrhea
- Fast or irregular heartbeat
- Severe fatigue or weakness

YESCARTA is a trademark of Kite Pharma.
© 2017 Kite Pharma | CCRC-00008 10/2017



2355 Utah Avenue, El Segundo, CA 90245



Patient Wallet Card

Carry this card with you at all times. SHOW THIS CARD if you go to the emergency room or see any physician.

Tell any healthcare provider that sees you that you are being treated with YESCARTA™.

Stay within close proximity (within 2 hours) of the location where you received your treatment for at least 4 weeks after getting YESCARTA™.

Important Information for Healthcare Providers

Name of treating oncologist:

Office phone:

After-hours phone:

Date of YESCARTA™ (axicabtagene ciloleucel) infusion:

- This patient has received YESCARTA™, which is a CD19-directed genetically modified autologous T-cell immunotherapy

- **YESCARTA™ can cause cytokine release syndrome (CRS) and neurologic toxicities, which may be fatal or life threatening. CRS may involve any organ system**



- **Contact the patient's oncologist immediately for further information**